arXiv:1008.1436v1 [math.NT] 9 Aug 2010

A family of generalized q-Genocchi numbers and polynomials

T. Kim[1], Byungje Lee[2], C. S. Ryoo[3]

[1] Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Korea
[2] Depart. of Wireless Communications Engineering, Kwangwoon University, Seoul 139-701, Korea
[3] Department of Mathematics, Hannam University, Daejeon 306-791, Korea

Abstract : In this paper we first consider the q-extension of the generating function for the higher-order generalized Genocchi numbers and polynomials attached to χ. The purpose of this paper is to present a systemic study of some families of higher-order generalized q-Genocchi numbers and polynomials attached to χ by using the generating function of those numbers and polynomials.

2000 Mathematics Subject Classification : 11B68, 11S40, 11S80

Key words : Genocchi numbers, Genocchi polynomials, higher-order generalized q-Genocchi numbers and polynomials

1. Introduction

As well known definition, the Genocchi polynomials are defined by

$$\left(\frac{2t}{e^t+1}\right)e^{xt}=e^{G(x)t}=\sum_{n=0}^{\infty}G_n(x)\frac{t^n}{n!},\quad |t|<\pi, \tag{1}$$

where we use the technical method's notation by replacing $G^n(x)$ by $G_n(x)$, symbolically, (see [3,11]). In the special case $x=0$, $G_n=G_n(0)$ are called the n-th Genocchi numbers. From the definition of Genocchi numbers, we note that $G_1=1, G_3=G_5=G_7=\cdots=0$, and even coefficients are given by $G_{2n}=2(1-2^{2n})B_{2n}=2nE_{2n-1}(0)$, (see [8]), where B_n is a Bernoulli number and $E_n(x)$ is an Euler polynomial. The first few Genocchi numbers for $2,4,6,\cdots$ are $-1,1,-3,17,-155,2073,\cdots$. The first few prime Genocchi numbers are given by $G_6=-3$ and $G_8=17$. It is known that there are no others prime Genocchi numbers with $n<10^5$. For a real or complex parameter α, the higher-order Genocchi polynomials are defined by

$$\left(\frac{2t}{e^t+1}\right)^{\alpha}e^{xt}=\sum_{n=0}^{\infty}G_n^{(\alpha)}(x)\frac{t^n}{n!},\ \text{(see [3, 4]).} \tag{2}$$

In the special case $x=0$, $G_n^{(\alpha)}=G_n^{(\alpha)}(0)$ are called the n-th Genocchi numbers of order α. From (1) and (2), we note that $G_n=G_n^{(1)}$. For $d\in\mathbb{N}$ with $d\equiv 1 \pmod 2$, let χ be the Dirichlet character with conductor d. It is known that the generalized Genocchi polynomials attached to χ are defined by

$$\left(\frac{2t\sum_{a=0}^{d-1}\chi(a)(-1)^a e^{at}}{e^{dt}+1}\right)e^{xt}=\sum_{n=0}^{\infty}G_{n,\chi}(x)\frac{t^n}{n!},\ \text{(see [3]).} \tag{3}$$

In the special case $x=0$, $G_{n,\chi}=G_{n,\chi}(0)$ are called the n-th generalized Genocchi numbers attached to χ (see [3, 4, 5, 6]).

For a real or complex parameter α, the generalized higher-order Genocchi polynomials attached to χ are also defined by

$$\left(\frac{2t\sum_{a=0}^{d-1}\chi(a)(-1)^a e^{at}}{e^{dt}+1}\right)^{\alpha}e^{xt}=\sum_{n=0}^{\infty}G_{n,\chi}^{(\alpha)}(x)\frac{t^n}{n!},\ \text{(see [13]).} \tag{4}$$

In the special case $x = 0$, $G_{n,\chi}^{(\alpha)} = G_{n,\chi}^{(\alpha)}(0)$ are called the n-th generalized Genocchi numbers attached to χ of order α (see [3, 4, 5, 6, 13, 14, 15]). From (3) and (4), we derive $G_{n,\chi} = G_{n,\chi}^{(1)}$.

Let us assume that $q \in \mathbb{C}$ with $|q| < 1$ as an indeterminate. Then we use the notation

$$[x]_q = \frac{1-q^x}{1-q}.$$

The q-factorial is defined by

$$[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$$

and the Gaussian binomial coefficient is also defined by

$$\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q}{[k]_q!}, \text{ (see [2, 5]).} \tag{5}$$

Note that

$$\lim_{q\to 1} \binom{n}{k}_q = \binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k!}.$$

It is known that

$$\binom{n+1}{k}_q = \binom{n}{k-1}_q + q^k \binom{n}{k}_q = q^{n-k} \binom{n}{k-1}_q + \binom{n}{k}_q, \text{ (see [2, 5]).}$$

The q-binomial formula are known that

$$(x-y)_q^n = (x-y)(x-qy)\cdots(x-q^{n-1}y) = \sum_{i=0}^{n} \binom{n}{i}_q q^{\binom{i}{2}} (-1)^i x^{n-i} y^i, \text{ (see [2, 9]),}$$

and

$$\frac{1}{(x-y)_q^n} = \frac{1}{(x-y)(x-qy)\cdots(x-q^{n-1}y)} = \sum_{l=0}^{\infty} \binom{n+l-1}{l}_q x^{n-l} y^l, \text{ (see [2, 9]).} \tag{6}$$

There is an unexpected connection with q-analysis and quantum groups, and thus with noncommutative geometry q-analysis is a sort of q-deformation of the ordinary analysis. Spherical functions on quantum groups are q-special functions. Recently, many authors have studied the q-extension in various area(see [1-15]). Govil and Gupta [2] has introduced a new type of q-integrated Meyer-König-Zeller-Durrmeyer operators and their results are closely related to study q-Bernstein polynomials and q-Genocchi polynomials, which are treated in this paper. In this paper, we first consider the q-extension of the generating function for the higher-order generalized Genocchi numbers and polynomials attached to χ. The purpose of this paper is to present a systemic study of some families of higher-order generalized q-Genocchi numbers and polynomials attached to χ by using the generating function of those numbers and polynomials.

2. Generalized q-Genocchi numbers and polynomials

For $r \in \mathbb{N}$, let us consider the q-extension of the generalized Genocchi polynomials of order r attached to χ as follows:

$$F_{q,\chi}^{(r)}(t,x) = 2^r t^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{j=1}^{r} \chi(m_j)\right) (-1)^{\sum_{j=1}^{r} m_j} e^{[x+m_1+\cdots+m_r]_q t} = \sum_{n=0}^{\infty} G_{n,\chi,q}^{(r)}(x) \frac{t^n}{n!}. \tag{7}$$

Note that

$$\lim_{q\to 1} F_{q,\chi}^{(r)}(t,x) = \left(\frac{2t\sum_{a=0}^{d-1} \chi(a)(-1)^a e^{at}}{e^{dt}+1}\right)^r e^{xt}.$$

By (7) and (4), we can see that $\lim_{q\to 1} G_{n,\chi,q}^{(r)}(x) = G_{n,\chi}^{(r)}(x)$. From (7), we note that

$$\begin{aligned}
&G_{0,\chi,q}^{(r)}(x) = G_{1,\chi,q}^{(r)}(x) = \cdots = G_{r-1,\chi,q}^{(r)}(x) = 0, \text{ and} \\
&\frac{G_{n+r,\chi,q}^{(r)}(x)}{\binom{n+r}{r} r!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{j=1}^{r} \chi(m_j) \right) (-1)^{\sum_{j=1}^{r} m_j} [x + m_1 + \cdots + m_r]_q^n .
\end{aligned}$$

In the special case $x = 0$, $G_{n,\chi,q}^{(r)} = G_{n,\chi,q}^{(r)}(0)$ are called the n-th generalized q-Genocchi numbers of order r attached to χ. Therefore, we obtain the following theorem.

Theorem 1. For $r \in \mathbb{N}$, we have

$$\frac{G_{n+r,\chi,q}^{(r)}}{\binom{n+r}{r} r!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} [m_1 + \cdots + m_r]_q^n .$$

Note that

$$\begin{aligned}
&2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} [m_1 + \cdots + m_r]_q^n \\
&= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) \frac{(-q^l)^{\sum_{i=1}^{r} a_i}}{(1+q^{ld})^r} .
\end{aligned}$$

Thus we obtain the following corollary.

Corollary 2. For $r \in \mathbb{N}$, we have

$$\begin{aligned}
\frac{G_{n+r,\chi,q}^{(r)}}{\binom{n+r}{r} r!} &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) \frac{(-q^l)^{\sum_{i=1}^{r} a_i}}{(1+q^{ld})^r} \\
&= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m} (-1)^m \sum_{a_1,\cdots,a_r=0}^{d-1} (-1)^{\sum_{i=1}^{r} a_i} \left(\prod_{i=1}^{r} \chi(a_i) \right) \left[\sum_{i=1}^{r} a_i + md \right]_q^n .
\end{aligned}$$

For $h \in \mathbb{Z}$ and $r \in \mathbb{N}$, we also consider the extended higher-order generalized (h,q)-Genocchi polynomials as follows:

$$\begin{aligned}
F_{q,\chi}^{(h,r)}(t,x) &= 2^r t^r \sum_{m_1,\cdots,m_r=0}^{\infty} q^{\sum_{j=1}^{r} (h-j) m_j} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} e^{[x + \sum_{j=1}^{r} m_j]_q t} \\
&= \sum_{n=0}^{\infty} G_{n,\chi,q}^{(h,r)}(x) \frac{t^n}{n!} .
\end{aligned} \tag{8}$$

From (8), we note that

$$\begin{aligned}
&G_{0,\chi,q}^{(h,r)}(x) = G_{1,\chi,q}^{(h,r)}(x) = \cdots = G_{r-1,\chi,q}^{(h,r)}(x) = 0, \text{ and} \\
&\frac{G_{n+r,\chi,q}^{(h,r)}(x)}{\binom{n+r}{r} r!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} q^{\sum_{j=1}^{r} (h-j) m_j} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} [x + m_1 + \cdots + m_r]_q^n \\
&= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} q^{lx} (-1)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) q^{\sum_{j=1}^{r} (h-j) a_j} (-1)^{a_1 + \cdots + a_r} q^{l(a_1 + \cdots + a_r)} \\
&\qquad \times \sum_{m_1,\cdots,m_r=0}^{\infty} (-1)^{m_1 + \cdots + m_r} q^{d(m_1 + \cdots + m_r) + d(\sum_{j=1}^{r} (h-j) m_j)} \\
&= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \frac{\binom{n}{l} q^{lx} (-1)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) q^{\sum_{j=1}^{r} (h-j) a_j} (-q^l)^{\sum_{i=1}^{r} a_i}}{(-q^{d(h-r+l)}; q)_r} ,
\end{aligned}$$

where $(-x;q)_r = (1+x)(1+xq)\cdots(1+xq^{r-1})$.

Therefore, we obtain the following theorem.

Theorem 3. For $h \in \mathbb{Z}, r \in \mathbb{N}$, we have

$$\frac{G^{(h,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r}r!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} q^{\sum_{j=1}^{r}(h-j)m_j}\left(\prod_{i=1}^{r}\chi(m_i)\right)(-1)^{\sum_{j=1}^{r}m_j}[x+m_1+\cdots+m_r]_q^n$$
$$= \frac{2^r}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-q^x)^l\sum_{a_1,\cdots,a_r=0}^{d-1}\left(\prod_{j=1}^{r}\chi(a_j)\right)q^{\sum_{j=1}^{r}(h-j)a_j}(-q^l)^{\sum_{i=1}^{r}a_i}}{(-q^{d(h-r+l)};q)_r},$$
$$\text{and}\quad G^{(h,r)}_{0,\chi,q}(x) = G^{(h,r)}_{1,\chi,q}(x) = \cdots = G^{(h,r)}_{r-1,\chi,q}(x) = 0.$$

Note that

$$\frac{1}{(-q^{d(h-r+l)};q)_r} = \frac{1}{(1+q^{d(h-r+l)})} = \sum_{m=0}^{\infty}\binom{m+r-1}{m}_q(-1)^m q^{d(h-r+l)m}. \tag{9}$$

By (9), we see that

$$\frac{1}{(1-q)^n}\sum_{l=0}^{n}\frac{\binom{n}{l}(-1)^l q^{l(x+\sum_{i=1}^{r}a_i)}}{(-q^{d(h-r+l)};q)_r}$$
$$= \sum_{m=0}^{\infty}\binom{m+r-1}{m}_q(-1)^m q^{d(h-r)m}\frac{1}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-1)^l q^{l(x+\sum_{i=1}^{r}a_i+dm)} \tag{10}$$
$$= \sum_{m=0}^{\infty}\binom{m+r-1}{m}_q(-1)^m q^{d(h-r)m}[x+\sum_{i=1}^{r}a_i+dm]_q^n.$$

By (9) and (10), we obtain the following corollary.

Corollary 4. For $h \in \mathbb{Z}, r \in \mathbb{N}$, we have

$$\frac{G^{(h,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r}r!}$$
$$= 2^r\sum_{m=0}^{\infty}\binom{m+r-1}{m}_q(-1)^m q^{d(h-r)m}\sum_{a_1,\cdots,a_r=0}^{d-1}\left(\prod_{j=1}^{r}\chi(a_j)\right)q^{\sum_{j=1}^{r}(h-j)a_j}[x+\sum_{i=1}^{r}a_i+dm]_q^n.$$

By (8), we can derive the following corollary.

Corollary 5. For $h \in \mathbb{Z}, r, d \in \mathbb{N}$ with $d \equiv 1(\mod 2)$, we have

$$q^{d(h-1)}\frac{G^{(h,r)}_{n+r,\chi,q}(x+d)}{\binom{n+r}{r}r!} + \frac{G^{(h,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r}r!} = 2\sum_{l=0}^{d-1}\chi(l)(-1)^l\frac{G^{(h-1,r-1)}_{n+r-1,\chi,q}}{\binom{n+r-1}{r-1}(r-1)!},$$

and

$$q^x\frac{G^{(h+1,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r}r!} = (q-1)\frac{G^{(h,r)}_{n+r+1,\chi,q}(x)}{\binom{n+r+1}{r}r!} + \frac{G^{(h,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r}r!}.$$

For $h = r$ in Theorem 3, we obtain the following corollary.

Corollary 6. For $r \in \mathbb{N}$, we have

$$\begin{aligned} & G^{(r,r)}_{n+r,\chi,q}(x) \\ &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-q^x)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) \frac{q^{\sum_{j=1}^{r}((r-j)a_j + la_j)}(-1)^{a_1+\cdots+a_r}}{(-q^{dl};q)_r} \\ &= 2^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-1)^m \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) q^{\sum_{j=1}^{r}(r-j)a_j} [x + \sum_{i=1}^{r} a_i + dm]_q^n. \end{aligned}$$

In particular,

$$\frac{G^{(r,r)}_{n+r,\chi,q^{-1}}(r-x)}{\binom{n+r}{r} r!} = (-1)^n q^{n+\binom{r}{2}} \frac{G^{(r,r)}_{n+r,\chi,q}(x)}{\binom{n+r}{r} r!}.$$

Let $x = r$ in Corollary 6. Then we have

$$\frac{G^{(r,r)}_{n+r,\chi,q^{-1}}}{\binom{n+r}{r} r!} = (-1)^n q^{n+\binom{r}{2}} \frac{G^{(r,r)}_{n+r,\chi,q}(r)}{\binom{n+r}{r} r!}.$$

Let $w_1, w_2, \cdots, w_r \in \mathbb{Q}_+$. Then we define Barnes' type generalized q-Genocchi polynomials attached to χ as follows:

$$\begin{aligned} F^{(r)}_{q,\chi}(t, x \mid w_1, w_2, \cdots, w_r) &= 2^r t^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{m_1+\cdots+m_r} e^{[x+\sum_{j=1}^{r} w_j m_j]_q t} \\ &= \sum_{n=0}^{\infty} G^{(r)}_{n,\chi,q}(x \mid w_1, w_2, \cdots, w_r) \frac{t^n}{n!}. \end{aligned} \tag{11}$$

By (11), we see that

$$\frac{G^{(r)}_{n+r,\chi,q}(x \mid w_1, \cdots, w_r)}{\binom{n+r}{r} r!} = 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} [x + \sum_{j=1}^{r} w_j m_j]_q^n.$$

It is easy to see that

$$\begin{aligned} & 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{m_1+\cdots+m_r} [x + \sum_{j=1}^{r} w_j m_j]_q^n \\ &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \frac{\binom{n}{l}(-q^x)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) (-1)^{\sum_{j=1}^{r} a_j} q^{l \sum_{i=1}^{r} w_i a_i}}{(1+q^{dlw_1}) \cdots (1+q^{dlw_r})}. \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 7. For $r \in \mathbb{N}, w_1, w_2, \cdots, w_r \in \mathbb{Q}_+$, we have

$$\begin{aligned} & \frac{G^{(r)}_{n+r,\chi,q}(x \mid w_1, w_2, \cdots, w_r)}{\binom{n+r}{r} r!} \\ &= 2^r \sum_{m_1,\cdots,m_r=0}^{\infty} \left(\prod_{i=1}^{r} \chi(m_i) \right) (-1)^{\sum_{j=1}^{r} m_j} [x + w_1 m_1 + \cdots + w_r m_r]_q^n \\ &= \frac{2^r}{(1-q)^n} \sum_{l=0}^{n} \frac{\binom{n}{l}(-q^x)^l \sum_{a_1,\cdots,a_r=0}^{d-1} \left(\prod_{j=1}^{r} \chi(a_j) \right) (-1)^{\sum_{j=1}^{r} a_j} q^{l \sum_{i=1}^{r} w_i a_i}}{(1+q^{dlw_1}) \cdots (1+q^{dlw_r})}. \end{aligned}$$

REFERENCES

1. I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, On the higher-order w-q-Genocchi numbers, Adv. Stud. Contemp. Math., 19(2009), 39-57.
2. N. K. Govil, V. Gupta, Convergence of q-Meyer-König-Zeller-Durrmeyer operators, Adv. Stud. Contemp. Math., 19(2009), 97-108.
3. L.-C. Jang, K.-W. Hwang, Y.-H. Kim, A note on (h,q)-Genocchi polynomials and numbers of higher order, Adv. Difference. Equ., 2010, Art ID 309480, 6pp.
4. L.-C. Jang, A study on the distribution of twisted q-Genocchi polynomials, Adv. Stud. Contemp. Math., 18(2009), 181-189.
5. T. Kim, Some identities for the Bernoulli, the Euler and the Genocchi numbers and polynomials, Adv. Stud. Contemp. Math., 20(2010), 23-28.
6. T. Kim, A note on the q-Genocchi numbers and polynomials, J. Inequal. Appl., 2007, Art ID 71452, 8pp.
7. T. Kim, On the multiple q-Genocchi and Euler numbers, Russ. J. Math. Phys., 15(2008), 481-486.
8. T. Kim, On the q-extension of Euler and Genocchi numbers, J. Math. Anal. Appl., 326(2007) 1458-1465.
9. T. Kim, Barnes type multiple q-zeta function and q-Euler polynomials, J. Phys. A: Math. Theor., 43(2010) 255201, 11pp.
10. T. Kim, Note on the Euler q-zeta functions, J. Number Theory, 129(2009), 1798-1804.
11. V. Kurt, A further symmetric relation on the analogue of the Apostol-Bernoulli and the analogue of the Apostol-Genocchi polynomials, Appl. Math. Sci., 3(2009), 2757-2764.
12. M. Cenkci, M. Can, V. Kurt, q-extension of Genocchi numbers, J. Korean Math. Soc., 43(2006), 183-198.
13. S.-H. Rim, S. J. Lee, E. J. Moon, J. H. Jin, On the q-Genocchi numbers and polynomials associated with q-zeta function, Proc. Jangjeon Math. Soc., 12(2009), 261-267.
14. S.-H. Rim, K. H. Park, E. J. Moon, On Genocchi numbers and polynomials, Abstr. Appl. Anal., Art ID 898471, 7pp.
15. C. S. Ryoo, Calculating zeros of the twisted Genocchi polynomials, Adv. Stud. Contemp. Math., 17(2008), 147-159.